



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## HYMENORUS OBSCURUS AS A PEST OF CITRUS (COL. CISTELIDAE)

J. R. WATSON

Occasionally one sees on the bark of citrus and other trees a dense colony composed of hundreds of little black beetles. The beetles are oval in shape, less than a quarter of an inch long and covered with grayish-brown hairs. In the late afternoon the beetles leave their resting place and go in search of food which consists mostly of lichens and other growths on the bark of the trees. But it seems that they may occasionally become pests. In July Mr. S. B. Jones of Orchid, Fla., sent in to the Experiment Station a number of these beetles with the statement that they had been feeding extensively on "June bloom" and other tender growth of his trees. In confinement they feed greedily on tender citrus foliage.

The writer has also recently caught these beetles eating out freshly inserted buds in a nursery. It would seem that this beetle must be included among the minor pests of a citrus tree.

This beetle should not be confused with the downy darkling beetle (*Epitragus tomentosus*), which it considerably resembles in shape and color. The latter is larger, never collects in colonies, and is one of the most beneficial insects in a citrus grove. Its food habits are very similar to those of lady beetles and in many groves it is much more abundant than even the twice-stabbed lady-beetle.

---

## A NEW CITRUS INSECT

A caterpillar recently found feeding on the leaves of a young grapefruit tree at Orlando, Florida, proved on rearing to maturity to be *Prodenia latifascia* Walker.

Altho this insect probably is of no economic importance as a pest, its presence seems to be a new record for citrus insects. Hence this note may be of interest.

The identification was made by Mr. Wm. Schaus of the National Museum, who says that *P. latifascia* Walker is essentially a tropical insect found from Mexico to Argentina, including Jamaica, Cuba, Haiti, St. Lucia. The only previous records of its breeding in the United States is one each from onions and alfalfa in Texas. Nothing is known of its host plants in other countries.